

## Collagen 4

## IN VITRO DIAGNOSTIC DATASHEET

INTENDED USE: IN VITRO DIAGNOSTIC USE

This product is intended for qualitative immunohistochemistry with normal and neoplastic formalin-fixed, paraffin-embedded tissue sections, to be viewed by light microscopy.

**DESCRIPTION:** Collagen is the main protein of connective tissue in animals and the most abundant protein in mammals, making up about 25% of the total protein content. Collagen IV is a major constituent of the basement membranes, along with laminins and enactins. It is composed of the alpha 1 IV chain and alpha 2 IV chain in a 2:1 ratio. It can form insoluble fibers with high tensile strength.

 CATALOG NO:
 PL747
 PL747-R7
 7 ML RTU 70 TEST

 PL747-R1
 1 ML RTU 10 TEST

 STAINING PATTERN:
 Cytoplasm
 PL747-1
 1 ML 1/100 1000 TEST

 PL747-0,1
 0,1 ML 1/100 100 TEST

POSITIVE CONTROL: Muscle, Lung

VOLUME: 7 ml Ready to Use (7 ml of antibody prediluted in 0.05mol/L Tris-HCl, pH 7.6 containing

stabilizing protein and 0.015mol/L sodium azide.)

**HOST:** Mouse

CLONE: CIV22

ANTIBODY CONCENTRATION: Not defined.

SPECIES REACTIVITY: Human. Others not tested.

EPITOPE: Not defined.

**MICROBIOLOGICAL STATE:** This product is not sterile.

**PRETREATMENT:** Staining of formalin-fixed tissue sections requires treating the tissue sections in boiling 10mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at room temperature for 20 min.

PRIMARY ANTIBODY INCUBATION TIME: 60 minutes at Room Temperature

STAINING TIPS: If the staining is too light, use lower dilution or longer time. If the staining is

too strong, check pretreatment, use higher dilution or shorter time.

STORAGE AND STABILITY: This product contains sodium azide and is stable for 24 months when stored

at 2-8°C. Do not use after expiration date indicated on label of the product. If reagent

is not stored as recommended, performance must be validated by the user.

TROUBLESHOOTING: Please contact Patolab Technical Support by e-mail (patolab@patolab.com.tr).



