

MSH-6

IN VITRO DIAGNOSTIC DATASHEET

INTENDED USE : IN VITRO DIAGNOSTIC USE

This product is intended for qualitative immunohistochemistry with normal and neoplastic formalin-fixed, paraffin-embedded tissue sections, to be viewed by light microscopy.

DESCRIPTION : MSH6 is a gene commonly associated with Hereditary Non-Polyposis Colorectal Cancer (HNPCC). HNPCC is an autosomal, dominantly inherited disease associated with marked increase in cancer susceptibility. It is characterized by a familial predisposition to early onset Colorectal Carcinoma and extra-colonic cancers of the gastrointestinal, urological and female reproductive tracts. HNPCC is reported to be the most common form of inherited Colorectal Cancer in the western world. MSH6 is a mismatch repair gene which is deficient in a high proportion of patients with microsatellite instability (MSI-H).

| | PL2119-R7 7 ML RTU 70 TEST PL2119-R1 1 ML RTU 10 TEST |
|---|--|
| STAINING PATTERN : Nuclear | PL2119-1 1 ML 1/100 1000 TEST |
| POSITIVE CONTROL : Colon Mucosa, Colon Carcinoma | PL2119-0,1 0,1 ML 1/100 100 TEST |
| VOLUME : 7 ml Ready to Use (7 ml of antibody prediluted in 0.05mol/L Tris-HCl, pH 7.6 containing stabilizing protein and 0.015mol/L sodium azide.) HOST : Mouse | |
| | |
| CLONE : EPR3945 | |
| ANTIBODY CONCENTRATION : 200ug/ml | |
| SPECIES REACTIVITY: Human, Rat, Mouse, Dog. Others-not known. | |
| EPITOPE : Not determined | |
| MICROBIOLOGICAL STATE : This product is not sterile. | |
| PRETREATMENT : Staining of formalin-fixed tissue sections requires treating the tissue sections in boiling 10mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at room temperature for 20 min. | |
| PRIMARY ANTIBODY INCUBATION TIME : 30 minutes at Room Temperature | |
| STAINING TIPS : If the staining is too light, use lower dilution too strong, check pretreatment, use higher | |
| STORAGE AND STABILITY : This product contains sodium azide and is stable for 24 months when stored at 2-8°C. Do not use after expiration date indicated on label of the product. If reagent is not stored as recommended, performance must be validated by the user. | |
| TROUBLESHOOTING : Please contact Patolab Technical S | Support by e-mail (patolab@patolab.com.tr). |

PatoLab Biyomedikal Cihazlar ve Sarf Malzemeler San. ve Tic. Ltd. Şti.patolab@patolab.com.trwww.patolab.com.tr

