

## Calretinin

## IN VITRO DIAGNOSTIC DATASHEET

INTENDED USE : IN VITRO DIAGNOSTIC USE

This product is intended for qualitative immunohistochemistry with normal and neoplastic formalin-fixed, paraffin-embedded tissue sections, to be viewed by light microscopy. it is Rabbit Monoclonal.

**DESCRIPTION :** Recognizes a protein named Calretinin. Calretinin is an intracellular calcium-binding protein belonging to the troponin C superfamily characterized by a structural mitif described as the EF-hand domain. The immunohistochemical detection of calretinin in developing cerebellum is restricted to the later stages indicated by weak staining from week 21 of gestation, in Purkinje and basket cells and in neurons of the dentate nucleus. The intensity of staining increases as the cerebellum matures. In tumors, calretinin has been detected in mesotheliomas and some pulmonary adenocarcinomas.

CATALOG NO: PL9113	PL9113-R7	7 ML RTU 70 TEST
	PL9113-R1	1 ML RTU 10 TEST
STAINING PATTERN : Cytoplasmic	PL9113-1	1 ML 1/200 2000 TEST
	PL9113-0,1	0,1 ML 1/200 200 TEST
POSITIVE CONTROL : Mesothelioma		
<b>VOLUME :</b> 7 ml Ready to Use ( 7 ml of antibody prediluted in 0.05mol/L Tris-HCl, pH 7.6 containing stabilizing protein and 0.015mol/L sodium azide. )		
HOST : Rabbit		
CLONE : SP13		
ANTIBODY CONCENTRATION : Not known		
SPECIES REACTIVITY: Human. Others not tested.		
EPITOPE : Not known		
MICROBIOLOGICAL STATE : This product is not sterile.		
<b>PRETREATMENT</b> : Staining of formalin-fixed tissue sections requires treating the tissue sections in boiling 10mM Citrate, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.		
PRIMARY ANTIBODY INCUBATION TIME: 30 minutes at Room Temperature		
<b>STAINING TIPS :</b> If the staining is too light, use lower dilution or longer time. If the staining is too strong, check pretreatment, use higher dilution or shorter time.		
<b>STORAGE AND STABILITY :</b> This product contains sodium azide and is stable for 24 months when stored at 2-8°C. Do not use after expiration date indicated on label of the product. If reagent is not stored as recommended, performance must be validated by the user.		
<b>TROUBLESHOOTING :</b> Please contact Patolab Technical Support by e-mail (patolab@patolab.com.tr).		

