

## **Uroplakin 3**

IN VITRO DIAGNOSTIC DATASHEET

INTENDED USE: IN VITRO DIAGNOSTIC USE

This product is intended for qualitative immunohistochemistry with normal and neoplastic formalin-fixed, paraffin-embedded tissue sections, to be viewed by light microscopy.

**DESCRIPTION:** The protein encoded by this gene is a member of the transmembrane 4 superfamily, also known as the tetraspanin family. Most of these members are cell-surface proteins that are characterized by the presence of four hydrophobic domains. The proteins mediate signal transduction events that play a role in the regulation of cell development, activation, growth and motility. This encoded protein is found in the asymmetrical unit membrane (AUM) where it can complex with other transmembrane 4 superfamily proteins.

CATALOG NO: PL2522 PL2522-R7 7 ML RTU 70 TEST

PL2522-R1 1 ML RTU 10 TEST

STAINING PATTERN: Cytoplasmic and membranous PL2522-1 1 ML 1/100 1000 TEST

PL2522-0,1 0,1 ML 1/100 100 TEST

POSITIVE CONTROL: Bladder or urothelial carcinoma

VOLUME: 7 ml Ready to Use (7 ml of antibody prediluted in 0.05mol/L Tris-HCl, pH 7.6 containing

stabilizing protein and 0.015mol/L sodium azide.)

**HOST:** Mouse

CLONE: ZM204

ANTIBODY CONCENTRATION: Not defined.

SPECIES REACTIVITY: Human. Others-not known.

**EPITOPE:** Not defined.

**MICROBIOLOGICAL STATE:** This product is not sterile.

**PRETREATMENT:** Staining of formalin-fixed tissue sections requires treating the tissue sections in boiling 10mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at room temperature for 20 min.

PRIMARY ANTIBODY INCUBATION TIME: 60 minutes at Room Temperature

**STAINING TIPS:** If the staining is too light, use lower dilution or longer time. If the staining is

too strong, check pretreatment, use higher dilution or shorter time.

**STORAGE AND STABILITY:** This product contains sodium azide and is stable for 24 months when stored

at 2-8°C. Do not use after expiration date indicated on label of the product. If reagent

is not stored as recommended, performance must be validated by the user.

**TROUBLESHOOTING:** Please contact Patolab Technical Support by e-mail ( patolab@patolab.com.tr ).



