

Glycophorin A

IN VITRO DIAGNOSTIC DATASHEET

INTENDED USE : IN VITRO DIAGNOSTIC USE

This product is intended for qualitative immunohistochemistry with normal and neoplastic formalin-fixed, paraffin-embedded tissue sections, to be viewed by light microscopy.

DESCRIPTION : Glycophorin A is a sialoglycoprotein present on red blood cells (RBC) and erythroid precursor cells. It has been shown that glycophorins acts as the receptor for Sandei virus and parvovirus.

CATALOG NO :	PL1843	PL1843-R7	7 ML RTU 70 TEST
		PL1843-R1	1 ML RTU 10 TEST
STAINING PATTERN :	Cell membrane	PL1843-1	1 ML 1/200 2000 TEST
		PL1843-0,1	0,1 ML 1/200 200 TEST

POSITIVE CONTROL : Human erythrocytes in placenta

VOLUME : 7 ml Ready to Use (7 ml of antibody prediluted in 0.05mol/L Tris-HCl, pH 7.6 containing stabilizing protein and 0.015mol/L sodium azide.)

HOST : Mouse

CLONE : JC159

ANTIBODY CONCENTRATION : 200ug/ml

SPECIES REACTIVITY : Human, Rat, and Pig and chicken. Others-not known

EPITOPE : Extracellular domain

MICROBIOLOGICAL STATE : This product is not sterile.

PRETREATMENT : No special pretreatment is required for the immunohistochemistry of formalin-fixed, paraffin-embedded tissues.

PRIMARY ANTIBODY INCUBATION TIME : 30 minutes at Room Temperature

STAINING TIPS : If the staining is too light, use lower dilution or longer time. If the staining is too strong, check pretreatment, use higher dilution or shorter time.

STORAGE AND STABILITY : This product contains sodium azide and is stable for 24 months when stored at 2-8°C. Do not use after expiration date indicated on label of the product. If reagent is not stored as recommended, performance must be validated by the user.

TROUBLESHOOTING : Please contact Patolab Technical Support by e-mail (patolab@patolab.com.tr).