

Collagen 4

INTENDED USE : IN VITRO DIAGNOSTIC USE

This product is intended for qualitative immunohistochemistry with normal and neoplastic formalin-fixed, paraffin-embedded tissue sections, to be viewed by light microscopy.

DESCRIPTION : Collagen is the main protein of connective tissue in animals and the most abundant protein in mammals, making up about 25% of the total protein content. Collagen IV is a major constituent of the basement membranes, along with laminins and enactins. It is composed of the alpha 1 IV chain and alpha 2 IV chain in a 2:1 ratio. It can form insoluble fibers with high tensile strength.

CATALOG NO :	PL747	PL747-R7	7 ML RTU 70 TEST
		PL747-R1	1 ML RTU 10 TEST
STAINING PATTERN :	Cytoplasm	PL747-1	1 ML 1/100 1000 TEST
		PL747-0,1	0,1 ML 1/100 100 TEST
POSITIVE CONTROL :	Muscle, Lung		

VOLUME : 7 ml Ready to Use (7 ml of antibody prediluted in 0.05mol/L Tris-HCl, pH 7.6 containing stabilizing protein and 0.015mol/L sodium azide.)

HOST : Mouse

CLONE : CIV22

ANTIBODY CONCENTRATION : Not defined.

SPECIES REACTIVITY : Human. Others-not known.

EPITOPE : Not defined.

MICROBIOLOGICAL STATE : This product is not sterile.

PRETREATMENT : Staining of formalin-fixed tissue sections requires treating the tissue sections in boiling 10mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at room temperature for 20 min.

PRIMARY ANTIBODY INCUBATION TIME : 60 minutes at Room Temperature

STAINING TIPS : If the staining is too light, use lower dilution or longer time. If the staining is too strong, check pretreatment, use higher dilution or shorter time.

STORAGE AND STABILITY : This product contains sodium azide and is stable for 24 months when stored at 2-8°C. Do not use after expiration date indicated on label of the product. If reagent is not stored as recommended, performance must be validated by the user.

TROUBLESHOOTING : Please contact Patolab Technical Support by e-mail (patolab@patolab.com.tr).

