

Desmin

IN VITRO DIAGNOSTIC DATASHEET

INTENDED USE: IN VITRO DIAGNOSTIC USE

This product is intended for qualitative immunohistochemistry with normal and neoplastic formalin-fixed, paraffin-embedded tissue sections, to be viewed by light microscopy.

DESCRIPTION: Desmin is an intermediate filament protein of both smooth and striated muscles. Antibody to desmin reacts with striated (skeletal and cardiac) as well as smooth muscle cells. In skeletal and cardiac muscles, the staining is confined to the Z-bands giving a characteristic striated appearance. Anti-desmin antibody is useful in identification of tumors of myogenic origin. It reacts with leiomyosarcomas (smooth muscle) as well as rhabdomyosarcomas (striated muscle).

CATALOG NO: PL376 PL376-R7 7 ML RTU 70 TEST

PL376-R1 1 ML RTU 10 TEST

STAINING PATTERN: Cytoplasmic PL376-1 1 ML 1/200 2000 TEST

PL376-0,1 0,1 ML 1/200 200 TEST

POSITIVE CONTROL: Muscle or Sarcoma

VOLUME: 7 ml Ready to Use (7 ml of antibody prediluted in 0.05mol/L Tris-HCl, pH 7.6 containing

stabilizing protein and 0.015mol/L sodium azide.)

HOST: Mouse

CLONE: D33

ANTIBODY CONCENTRATION: Not known

SPECIES REACTIVITY: Human, Baboon, Monkey, Cow, Cat, Dog, Hamster, Rat, Chicken. Others-not known

EPITOPE: Not determined

MICROBIOLOGICAL STATE: This product is not sterile.

PRETREATMENT: No special pretreatment is required for immuno-histochemistry of formalin-fixed tissues.

PRIMARY ANTIBODY INCUBATION TIME: 30 minutes at Room Temperature

STAINING TIPS: If the staining is too light, use lower dilution or longer time. If the staining is

too strong, check pretreatment, use higher dilution or shorter time.

STORAGE AND STABILITY: This product contains sodium azide and is stable for 24 months when stored

at 2-8°C. Do not use after expiration date indicated on label of the product. If reagent

is not stored as recommended, performance must be validated by the user.

TROUBLESHOOTING: Please contact Patolab Technical Support by e-mail (patolab@patolab.com.tr).



