# Pat Lab<sup>®</sup>

# **AEC SUBSTRATE & CHROMOGEN KIT**

# **INTENDED USE:**

For In Vitro Diagnostic Use

| Catalog No | Volume               |  |
|------------|----------------------|--|
| PL-060-HA  | 60 ml ( 600 slides)  |  |
| PL-125-HA  | 125 ml (1250 slides) |  |

#### DESCRIPTION

3-Amino-9-Ethylcarbazole (AEC) is a widely used chromogen for immunohistochemical staining and immunoblotting. AEC produces a red end product that is soluble in alcohol and must be used with an aqueous counterstain and mounting media. This product is available in a two-component form consisting of a liquid, refrigerator-stable AEC Chromogen and AEC Substrate. Each component is individually stable for 18 months from the date of manufacture.

| <u>COMPONENTS</u> | <u>60 ML</u><br>PL-060-HA | <u>125 ML</u><br>PL-125-HA |
|-------------------|---------------------------|----------------------------|
| AEC Chromogen     | PL-006-HAC                | PL-012-HAC                 |
| AEC Substrate     | PL-060-HAS                | PL-125-HAS                 |

#### **WARNINGS & PRECAUTIONS**

Refer to MSDS.

# **STORAGE & SHELF LIFE**

Store at 2-8°C. Each component is stable for 24 months.

# **MICROBIOLOGICAL STATE**

Product(s) not sterile.

# MATERIALS REQUIRED BUT NOT PROVIDED

N/A

#### **SPECIMEN & REAGENT PREPARATION**

Refer to Procedure.

# PROCEDURE

- 1. Add 20 ul of AEC Chromogen to 1 ml of AEC Substrate and mix well.
- 2. Apply mixture to tissue section within 20 minutes of mixing.
- 3. Incubate tissue section for 5-20 minutes.
- 4. Rinse tissue, counterstain, and coverslip.

#### **Troubleshooting**

Please contact PatoLab Technical Support by e-mail (patolab@patolab.com.tr).

PatoLab Biyomedikal Cihazlar ve Sarf Malzemeler San. ve Tic. Ltd. Şti.patolab@patolab.com.trwww.patolab.com.tr

