

Keratin 7

IN VITRO DIAGNOSTIC DATASHEET

INTENDED USE: IN VITRO DIAGNOSTIC USE

This product is intended for qualitative immunohistochemistry with normal and neoplastic formalin-fixed, paraffin-embedded tissue sections, to be viewed by light microscopy.

DESCRIPTION: Cytokeratin 7 is a basic cytokeratin which is found in most glandular and transitional epithelia but not in the stratified squamous epithelia. Keratin 7 is expressed in the epithelial cells of ovary, lung, and breast but not of colon, prostate, or gastrointestinal tract. Antibody to cytokeratin is useful in distinguishing ovarian carcinomas (keratin 7+) from colon carcinomas (keratin 7-).

 CATALOG NO:
 PL1352
 PL1352-R7
 7 ML RTU 70 TEST

 PL1352-R1
 1 ML RTU 10 TEST

 STAINING PATTERN:
 Cytoplasmic
 PL1352-1
 1 ML 1/100 1000 TEST

 PL1352-0,1
 0,1 ML 1/100 100 TEST

POSITIVE CONTROL: Ovarian, Breast Carcinoma

VOLUME: 7 ml Ready to Use (7 ml of antibody prediluted in 0.05mol/L Tris-HCl, pH 7.6 containing

stabilizing protein and 0.015mol/L sodium azide.)

HOST: Mouse

CLONE: OV-TL 12/30

ANTIBODY CONCENTRATION: 200ug/ml

SPECIES REACTIVITY: Human. Does not react with rat. Others-not known

EPITOPE: Not determined

MICROBIOLOGICAL STATE: This product is not sterile.

PRETREATMENT: Staining of formalin-fixed tissue sections requires treating the tissue sections in boiling 10mM Citrate, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.

PRIMARY ANTIBODY INCUBATION TIME: 30 minutes at Room Temperature

STAINING TIPS: If the staining is too light, use lower dilution or longer time. If the staining is

too strong, check pretreatment, use higher dilution or shorter time.

STORAGE AND STABILITY: This product contains sodium azide and is stable for 24 months when stored

at 2-8°C. Do not use after expiration date indicated on label of the product. If reagent is not stored as recommended, performance must be validated by the user.

TROUBLESHOOTING: Please contact Patolab Technical Support by e-mail (patolab@patolab.com.tr).



