

## Glypican-3

IN VITRO DIAGNOSTIC DATASHEET

**INTENDED USE :** IN VITRO DIAGNOSTIC USE

This product is intended for qualitative immunohistochemistry with normal and neoplastic formalin-fixed, paraffin-embedded tissue sections, to be viewed by light microscopy.

**DESCRIPTION :**

Glypican-3 (GPC3) is a member of the glypican family of glycosyl phosphatidylinositol-anchored cell-surface heparan sulfate proteoglycans. The 1G12 monoclonal antibody has been used to assess GPC3 expression in malignant and nonmalignant liver tissue samples by immunohistochemistry in formalin-fixed and paraffin-embedded tissue.

|                           |                            |            |                       |
|---------------------------|----------------------------|------------|-----------------------|
| <b>CATALOG NO :</b>       | PL1939                     | PL1939-R7  | 7 ML RTU 70 TEST      |
|                           |                            | PL1939-R1  | 1 ML RTU 10 TEST      |
| <b>STAINING PATTERN :</b> | Cytoplasmic and membranous | PL1939-1   | 1 ML 1/100 1000 TEST  |
|                           |                            | PL1939-0,1 | 0,1 ML 1/100 100 TEST |

**POSITIVE CONTROL :** Hepatocellular carcinoma

**VOLUME :** 7 ml Ready to Use ( 7 ml of antibody prediluted in 0.05mol/L Tris-HCl, pH 7.6 containing stabilizing protein and 0.015mol/L sodium azide. )

**HOST :** Mouse

**CLONE :** 1G12

**ANTIBODY CONCENTRATION :** Not defined.

**SPECIES REACTIVITY :** Human. Others not tested.

**EPITOPE :** Not defined.

**MICROBIOLOGICAL STATE :** This product is not sterile.

**PRETREATMENT :** Staining of formalin-fixed tissue sections requires treating the tissue sections in boiling 10mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at room temperature for 20 min.

**PRIMARY ANTIBODY INCUBATION TIME :** 60 minutes at Room Temperature

**STAINING TIPS :** If the staining is too light, use lower dilution or longer time. If the staining is too strong, check pretreatment, use higher dilution or shorter time.

**STORAGE AND STABILITY :** This product contains sodium azide and is stable for 24 months when stored at 2-8°C. Do not use after expiration date indicated on label of the product. If reagent is not stored as recommended, performance must be validated by the user.

**TROUBLESHOOTING :** Please contact Patolab Technical Support by e-mail ( patolab@patolab.com.tr ).